

The emergence of AdS_2 from quantum fluctuations

J. Ambjørn^{a,c} R. Janik^b, W. Westra^c and S. Zohren^d

^a The Niels Bohr Institute, Copenhagen University
 Blegdamsvej 17, DK-2100 Copenhagen Ø, Denmark.
 email: ambjorn@nbi.dk

^b Institute of Physics, Jagellonian University,
 ul. Reymonta 4, 30-059 Krakow, Poland.
 email: ufrjanik@if.uj.edu.pl

^c Institute for Theoretical Physics, Utrecht University,
 Leuvenlaan 4, NL-3584 CE Utrecht, The Netherlands.
 email: w.westra@phys.uu.nl, ambjorn@phys.uu.nl

^d Blackett Laboratory, Imperial College,
 London SW7 2AZ, United Kingdom.
 email: stefan.zohren@imperial.ac.uk

Abstract

We have shown how the quantization of two-dimensional quantum gravity with an action which contains only a positive cosmological constant and boundary cosmological constants leads to the emergence of a spacetime which can be described as a constant negative curvature spacetime with superimposed quantum fluctuations.

Talk given by W. Westra at the Eleventh Marcel Grossmann Meeting on General Relativity at the Freie U. Berlin, July 23 - 29, 2006.

1 Introduction

The causal dynamical triangulations approach to quantum gravity (CDT) is an attempt to define the gravitational path integral in a background independent and nonperturbative manner. As in the case of Euclidean dynamical triangulation approaches (DT), CDT provides a regularization of the path integral through a sum over piecewise linear geometries where the edge length of the individual building blocks serves as an ultraviolet cutoff. However, in contrast to DT a global time foliation is imposed on each individual history in the path integral.

The method of CDT was first applied to two-dimensional quantum gravity where the model was shown to be analytically solvable [1]. Although two-dimensional quantum gravity does not have any propagating degrees of freedom it is a fertile playground to study certain aspects of diffeomorphism invariant theories. Among the issues that have been addressed within the two-dimensional framework are the inclusion of a sum over topologies [2] and the emergence of a background geometry purely from quantum fluctuations [3] where the latter is the subject of this article.

One of the most natural quantities to study in CDT is the loop-loop-propagator which is the amplitude for a transition from a spacelike loop with boundary cosmological constant X to a loop with boundary cosmological constant Y in time T ,

$$G_\Lambda(X, Y; T) = \int \mathcal{D}[g] e^{-S[g]}, \quad S[g] = \Lambda \int d^2x \sqrt{g(x, t)} + X \oint dl_1 + Y \oint dl_2, \quad (1)$$

where the action only includes bulk and boundary cosmological constants, since the curvature term in the Einstein-Hilbert action is trivial in 2D.

Evaluating the path integral using the CDT regularization and taking the continuum limit yields the following [1]

$$G_\Lambda(X, Y; T) = \frac{\bar{X}^2(T, X) - \Lambda}{X^2 - \Lambda} \frac{1}{\bar{X}(T, X) + Y}, \quad (2)$$

where $\bar{X}(T, X)$ is the solution of

$$\frac{d\bar{X}}{dT} = -(\bar{X}^2 - \Lambda), \quad \bar{X}(0, X) = X. \quad (3)$$

2 The emergence of AdS_2

To determine the background geometry of the 1+1 dimensional universe we calculate the average spatial length at proper time $t \in [0, T]$

$$\langle L(t) \rangle_{X, Y, T} = \frac{1}{G_\Lambda(X, Y; T)} \int_0^\infty dL G_\Lambda(X, L; t) L G_\Lambda(L, Y; T - t). \quad (4)$$

Evaluating the average length at the boundary $t = T$ and taking the limit $T \rightarrow \infty$ gives

$$\lim_{T \rightarrow \infty} \langle L(T) \rangle_{X,Y,T} = \frac{1}{Y + \sqrt{\Lambda}}. \quad (5)$$

Interestingly, one observes that there is a special value $Y = -\sqrt{\Lambda}$ of the boundary cosmological constant for which the boundary length diverges and the geometry becomes non-compact. Using this critical value for the boundary cosmological constant Y one can obtain the boundary length for finite T

$$L^c(T) = \langle L(T) \rangle_{X,Y=-\sqrt{\Lambda},T} = \frac{1}{\sqrt{\Lambda}} \frac{1}{\coth \sqrt{\Lambda} T - 1}. \quad (6)$$

Instead of using boundary cosmological constants one can also fix the spatial length of the boundaries. Using the Laplace transformed propagator $G_\Lambda(L_1, L_2; T)$ we can evaluate the average spatial length $\langle L(t) \rangle_{L_1, L_2, T}$ for fixed lengths L_1 and L_2 of the boundary loops.

In the following we want to investigate the quantum geometry in the case where it becomes non-compact. Therefore we set the boundary length at $t = T$ to the critical value $L^c(T)$ as defined in (6) and for simplicity we shrink the spatial geometry at $t = 0$ to a point. In the limit $T \rightarrow \infty$ one obtains the average length of the spatial geometry at proper time $t \in [0, T]$

$$\langle L(t) \rangle \equiv \lim_{T \rightarrow \infty} \langle L(t) \rangle_{L_1=0, L_2=L^c(T), T} = \frac{1}{\sqrt{\Lambda}} \sinh(2\sqrt{\Lambda}t). \quad (7)$$

Due to the fact that L and T are defined from the continuum limit of a simplicial geometry there is a relative constant of proportionality that can only be fixed by comparing with continuum calculations [4] yielding $L_{cont}(t) = \pi \langle L(t) \rangle$. From this result the metric for the background geometry is readily obtained,

$$ds^2 = dt^2 + \frac{L_{cont}^2}{4\pi^2} d\theta^2 = dt^2 + \frac{\sinh^2(2\sqrt{\Lambda}t)}{4\Lambda} d\theta^2. \quad (8)$$

This is nothing but the metric of the Poincaré disc which can be seen as a Wick rotated version of AdS_2 with constant negative curvature $R = -8\Lambda$.

To better understand the quantum nature of the geometry it is useful to compute the fluctuations of the spatial length. From expressions analogous to (4) one can determine the relative fluctuations

$$\frac{\Delta L(t)}{\langle L(t) \rangle} \equiv \frac{\sqrt{\langle L^2(t) \rangle - \langle L(t) \rangle^2}}{\langle L(t) \rangle} \sim e^{-\sqrt{\Lambda}t}. \quad (9)$$

Surprisingly, the fluctuations of the spatial geometry become exponentially small for $t \gg \Lambda^{-1/2}$. Concluding from Eqs. (8) and (9), one can view the quantum geometry as a version of Wick rotated AdS_2 dressed with small quantum fluctuations.

3 Discussion

We have shown that in 2D quantum gravity defined through CDT there is a transition from compact geometry to non-compact AdS_2 -like geometry for a special value of the boundary cosmological constant. This phenomenon is similar to the Euclidean case where non-compact ZZ-branes appear in a transition from compact 2D geometries in Liouville quantum gravity [5]. A surprising feature of the CDT result is that the fluctuations become exponentially small which enables us to interpret the emerging AdS_2 spacetime as a genuine semiclassical background. It is interesting that similar results have been reported in four-dimensional CDT where numerical simulations indicate the emergence of a semi-classical background from a nonperturbative and background-independent path integral [6].

Acknowledgments

All authors acknowledge support by ENRAGE (European Network on Random Geometry), a Marie Curie Research Training Network in the European Community's Sixth Framework Programme, network contract MRTN-CT-2004-005616.

References

- [1] J. Ambjørn and R. Loll, *Nucl. Phys. B* **536**, 407 (1998) [hep-th/9805108].
- [2] R. Loll, W. Westra and S. Zohren, *Nucl. Phys. B* **751**, 419 (2006) [hep-th/0507012].
- [3] J. Ambjørn, R. Janik, W. Westra and S. Zohren, *Phys. Lett. B* **641**, 94 (2006) [gr-qc/0607013].
- [4] R. Nakayama, *Phys. Lett. B* **325**, 347 (1994) [hep-th/9312158].
- [5] J. Ambjørn, S. Arianos, J. A. Gesser and S. Kawamoto, *Phys. Lett. B* **599**, 306 (2004) [hep-th/0406108].
- [6] J. Ambjørn, J. Jurkiewicz and R. Loll, *Phys. Rev. Lett.* **93**, 131301 (2004) [arXiv:hep-th/0404156], *Phys. Lett. B* **607**, 205 (2005) [hep-th/0411152], *Phys. Rev.* **D72**, p. 064014 (2005) [hep-th/0505154].